

SEQUENCES OF ANALYTIC DISKS

EVGENY A. POLETSKY

ABSTRACT. The subject considered in this paper has, at least, three points of interest. Suppose that we have a sequence of one-dimensional analytic varieties in a domain in $\mathbb{C}^n$. The cluster of this sequence consists from all points in the domains such that every neighbourhood of such points intersects with infinitely many different varieties. The first question is: what analytic properties does the cluster inherit from varieties? We give a sufficient criterion when the cluster contains an analytic disk, but it follows from examples of Stolzenberg and Wermer that, in general, clusters can contain no analytic disks. So we study algebras of continuous function on clusters, which can be approximated by holomorphic functions or polynomials, and show that this algebras possess some analytic properties in all but explicitly pathological and uninteresting cases.

Secondly, we apply and results about clusters to polynomial hulls and maximal functions, finding remnants of analytic structures there too.

And, finally, due to more and more frequent appearances of analytic disks as tools in complex analysis, it seems to be interesting to look at their sequences to establish terminology, basic notation and properties.

§0. INTRODUCTION

Let us consider the following example.

Let A_j be a sequence of one-dimensional irreducible analytic subvarieties of a bounded domain D in $\mathbb{C}^n$. *The cluster* of this sequence is the set A of all points in D such that every ball, centered at a point $z \in A$, has non-empty intersections with infinitely many different varieties A_j . The question is: Does A remember its analytic origin? And if it does, what kind of analytic geometry it inherits?

By the uniformization theorem for each j and for each point $z \in A_j$ there is a holomorphic mapping f_j of the unit disk U into D such that $f_j(0) = z$ and $f_j(U) = A_j$.

1991 *Mathematics Subject Classification*. Primary: 32C25; secondary: 32E20, 32F05.

Key words and phrases. Sequences of analytic functions, polynomial hulls, maximal functions.

Partially supported by NSF grant #DMS-9101826

Moreover, all asymptotic values of f_j belong to the boundary ∂D of D . It is more convenient to consider sequences of functions f_j instead of sequences of varieties A_j because we can use the greater power of complex analysis. Also this replacement brings into the consideration other important cases which we are going to discuss now.

Let X be a compact in $\mathbb{C}^n$ and $\hat{X}$ is its polynomial hull. If z is a point of $\hat{X}$ then (see Th. 5.1 below) there is a sequence of holomorphic mappings $f_j : U \rightarrow \mathbb{C}^n$ such that $f_j(0) = z$ and boundary values of f_j belong to an arbitrary fixed neighbourhood V of X everywhere on the unit circle S except of a set of measure $\varepsilon_j \rightarrow 0$. Moreover, the cluster of the sequence $f_j(U)$ belongs to $\hat{X}$. Has the cluster an analytic structure in some sense?

The same question arises when one studies maximal function on the domain D in the pluripotential theory. It can be proved that values of these function can be obtained as solutions of variational problems for some functional Φ on the set of all holomorphic mappings of U into D . If $u(z)$ is the extremal value of Φ then we can find the sequence of holomorphic mappings f_j of U such that $f_j(0) = z$ and $\Phi(f_j) \rightarrow u$. Again, as before, boundary values of f_j lie near ∂D except of the set of a small measure. And, again, it is interesting to know how the maximal function behaves on the cluster of $f_j(U)$.

The question about the existence of analytic structure is, of course, ambiguous. To make it more specific we can ask whether the cluster has an analytic disk inside or not? This was the first question being studied, and Stolzenberg [12] and Wermer [14] provided examples of clusters which do not contain any non-trivial analytic disks. These are impressive examples which got titles of "clusters without analytic structure". We give several sufficient conditions for a cluster to contain an analytic disk. But one of main goals of this paper is a generalization of a result of Goldmann. In [6] Goldmann discovered that the Wermer's example has the uniqueness property: if a continuous function can be approximated by polynomials on this cluster and is equal to zero on some open set then it is equal to zero everywhere. So some analyticity stays in the cluster. We show in this paper that the uniqueness property holds in many quite general cases.

We also apply our results to maximal functions. Bedford and Kalka proved in [1] that if a such function is smooth then through every point one can draw an analytic disk such that the restriction of the function to this disk is harmonic. For continuous functions the result does not hold in this form, but if we replace analytic disks by clusters then everything is true.

There is one more application of introduced notions. Let $\{f_j : U \rightarrow \mathbb{C}^n\}$ be a uniformly bounded sequence of holomorphic mappings with weakly converging push-forward measures $(f_j)_*m = \mu(dz, f_j)$ of the measure $m = d\theta/2\pi$ on the unit circle. The weak limit of such measures is, evidently, a Jensen measure. Conversely, as it was proved in [3], every Jensen measure is such a limit. We do not explore this observation here concentrating on geometric properties of clusters.

And another goal of this paper is to develop the nomenclature for this kind of species, establish basic properties of clusters, and to show that they have quite ample geometry.

I am very grateful to Norm Levenberg for useful discussions.

§1. HOLOMORPHIC MEASURES

We shall denote by $B^n(a, r)$ and $S^n(a, r)$ the ball and the sphere of radius r with center a lying in the complex n -dimensional space $\mathbb{C}^n$. We shall omit a and r when $a = 0$ or $r = 1$ and the index n when this does not lead to a misunderstanding. We write U for B^1 and S for S^1 . A standard Lebesgue measure in $\mathbb{C}^n$ will be denoted by $m(dz)$.

If E is a set in the closure $\overline{D}$ a domain $D \subset \mathbb{C}$ then we define its harmonic measure in the following way. If $E \subset \overline{D}$ is an open set then the *harmonic measure* $\omega(z, E, D)$ of the set E with respect to D is equal to the infimum of all positive superharmonic functions on D which are continuous and greater than 1 on E . If F is an arbitrary set in D then we define the harmonic measure $\omega(z, F, D)$ to be equal to $\inf \omega(z, E, D)$, where the infimum is taken over all open sets E containing F . This function may be not upper semicontinuous, so we take its regularization

$$\overline{\lim}_{w \rightarrow z} \omega(z, F, D),$$

which is superharmonic and will be denoted also by $\omega(z, F, D)$.

Let f be a bounded holomorphic mapping of the unit disk into $\mathbb{C}^n$. Since f has radial limit values almost everywhere on S , we may consider f as a measurable mapping of the closed unit disk $\overline{U}$ and define for a Borel set $E \subset \mathbb{C}^n$ a set function

$$\mu(E, f) = \frac{1}{2\pi} m(f^{-1}(E) \cap S) = \omega(0, f^{-1}(E) \cap S, U).$$

This is a regular Borel measure on $\mathbb{C}^n$ and, if f is not constant, $\mu(dz, f)$ has no atoms, i.e. the measure μ of each point in $\mathbb{C}^n$ is zero. Let X be the support of μ . Evidently,

$f(U)$ belongs to the polynomial hull $\hat{X}$ of X , but $f(\overline{U})$ can be less than $\hat{X}$ as the example of Wermer in [13] shows. The point $z_f = f(0)$ is uniquely determined as a point such that

$$\int p(z) \mu(dz, f) = p(z_f)$$

for every polynomial p on $\mathbb{C}^n$. In particular,

$$\int z \mu(dz, f) = z_f.$$

We shall call the measure $\mu(dz, f)$ by a *holomorphic measure*, the point z_f by *the center of the holomorphic measure* $\mu(dz, f)$. An *analytic disk*, corresponding to this holomorphic measure, is the image $f(U)$ of f .

Since the author could not find any references in the literature concerning such measures, we shall describe some of their properties.

If $f = (f_1, f_2, \dots, f_n)$ then we may consider measures $\mu(dz, f_j)$ on $\mathbb{C}$ and, evidently, for a set E , lying in the j -th coordinate complex plane, we have $\mu(E, f_j) = \mu(E \times \mathbb{C}^{n-1}, f)$. Thus, the equality $\mu(dz, f) = \mu(dz, g)$ implies that $\mu(dz, f_j) = \mu(dz, g_j)$. The converse statement is, in general, false.

If analytic disk $f(U)$ is given, we may consider the mapping f as a parameterization of $f(U)$. We have the natural method of changing parameter by composing f with a mapping $h : U \rightarrow U$. Such a reparameterization does not change the center of the analytic disk if $h(0) = 0$ and does not change the holomorphic measure if and only if h is inner, as the following theorem shows.

Theorem 1.1. *For a holomorphic function $h : U \rightarrow U$ with $h(0) = 0$ measures $\mu(dz, f)$ and $\mu(dz, f \circ h)$ are equal if and only if h is an inner function, i.e. $|h| = 1$ a.e. on S .*

Proof. If E is a Borel set in $\mathbb{C}^n$ then the measure $\mu(E, f)$ is equal to the measure of the set $E_1 = f^{-1}(E) \cap S$. For the mapping $g = f \circ h$ the measure $\mu(E, g)$ is equal to the measure of the set $E_2 = g^{-1}(E) \cap S = h^{-1}(E_1)$. Since inner functions, mapping the origin into the origin, preserve measures on S (see [10, 19.3.4]), $m(E_2) = m(E_1)$.¹

Conversely, for a holomorphic mapping h of U into U , we consider the set A in S , where values of radial limits of h belongs to U . If $m(A) = 0$ then h is inner, so we may assume that $m(A) > 0$. Let A_* be the image of A under the mapping $g = f \circ h$. We

¹Prof. W. Rudin told me about this reference.

denote by A_1 the set $f^{-1}(A_*) \cap S$ and by A_2 the set $g^{-1}(A_*) \cap S$. Evidently, $A \subset A_2$ and we denote by A' the set $A_2 \setminus A$. Then

$$m(A_1) = m(A_2) = m(A) + m(A').$$

Let u be the harmonic function equal to 1 on A_1 and to zero on $S \setminus A_1$ and let $v = u \circ h$. Then $u(0) = v(0) = m(A_k)$, $k = 1, 2$. But the function v is equal to 0 on $S \setminus (A \cup A')$, to 1 on A' and is less than 1 on A . Therefore,

$$m(A_2) = m(A) + m(A') > v(0) = m(A_1). \quad \square$$

Holomorphic measures are related to the pluripotential theory as the following simple reasoning shows. A set $E \in \mathbb{C}^n$ is the set of *universal holomorphic measure zero* if for every nonconstant holomorphic mapping $f : U \rightarrow \mathbb{C}^n$ the measure $\mu(E, f) = 0$. Such sets exist, for example, all sets, containing only one point, are of universal holomorphic measure zero. In 20-th Lusin introduced the notion of P -set. A P -set E is a set in $\mathbb{R}^n$ such that $\mu(E) = 0$ for every finite Borel regular measure without atoms. There are non-trivial examples of P -sets.

Theorem 1.2.

- (1) *Every P -set in $\mathbb{C}^n$ is a set of universal holomorphic measure zero.*
- (2) *Every set of universal holomorphic measure zero in $\mathbb{C}^n$ is pluripolar.*
- (3) *A Borel set E in the complex plane $\mathbb{C}$ is a set of universal holomorphic measure zero if and only if it is polar.*

Proof. 1) is trivial.

Part 2) follows from Corollary 2.1.2 of [9].

Part 3) follows from part 2) and Theorem 2.16 of [4], claiming that polar sets has holomorphic measure zero for every non-constant bounded mapping $f : U \rightarrow \mathbb{C}$. $\square$

In $\mathbb{C}^n$ the latter statement does not hold. For example, take $E = \{(z, w) : ||(z, w)|| < 1, w = 0\}$, which is pluri-polar, but $\mu(E, f) \neq 0$ for $f(\zeta) = (\zeta, 0)$. It is interesting to describe all sets of universal holomorphic measure 0 in $\mathbb{C}^n$. Evidently,

Theorem 1.3. *Let A be a smooth complex analytic curve in a domain $D \subset \mathbb{C}^n$. A set $E \subset A$ is a set of universal holomorphic measure zero if and only if E is polar on A .*

Proof. Follows from part 3) of Theorem 1.2 $\square$

Of course, we must be aware that a set E can belong to supports of two holomorphic measures $\mu(dz, f)$ and $\mu(dz, g)$, and $\mu(E, f) \neq 0$ but $\mu(dz, g) = 0$.

Example 1.1. Let E be a closed set on $[-1, 1]$ with linear measure equal to zero but non-polar as a set in $\mathbb{C}$. Let f be a conformal mapping of U onto the upper half of U and let g be the universal covering of $U \setminus E$. Then $\mu(E, f) = 0$ and $\mu(E, g) > 0$.

It is interesting to find out when two holomorphic have equal holomorphic measures. It is easy to construct an example of two holomorphic mappings which measures are absolutely continuous with respect to each other, but not equal.

§2. JENSEN MEASURES.

A *Jensen measure* on $\mathbb{C}^n$ with barycenter $z_0 \in \mathbb{C}^n$ is a regular Borel measure μ with compact support such that $\mu(\mathbb{C}^n) = 1$ and for every plurisubharmonic function u on $\mathbb{C}^n$

$$u(z_0) \leq \int u(z) \mu(dz).$$

It is clear that every holomorphic measure $\mu(dz, f)$ is a Jensen measure with barycenter z_f .

Let $L = \{f_j\}$ be a sequence of uniformly bounded holomorphic mappings of U with weakly converging holomorphic measures $\mu(dz, f_j)$, i.e. for every continuous function $\varphi(z)$ on $\mathbb{C}^n$ we have

$$\lim_{j \rightarrow \infty} \int \varphi(z) \mu(dz, f_j) = \int \varphi(z) \mu(dz)$$

where μ is a regular Borel measure on $\mathbb{C}^n$. Then centers z_{f_j} converge to a point z_0 and the measure $\mu = \mu(dz, L)$ is a Jensen measure with barycenter z_0 . The following theorem, claiming that every Jensen measure on $\mathbb{C}^n$ can be obtained as a weak limit of holomorphic measures, was proved, basically, in [3]. Since we could not find the theorem in the form we need in [3] and for the readers' convenience we shall give another proof of this result.

Theorem 2.1. *Let D be a domain in $\mathbb{C}^n$ and let μ be a Jensen measure with barycenter $z_0 \in D$ and support compactly belonging to D . Then there is a sequence $L = \{f_j\}$ of holomorphic mappings of U into D with holomorphic measures $\mu(dz, f_j)$ weakly converging to μ .*

Proof. Let $z_0 = 0$. Following [3] we shall prove that the set $\mathbf{P}$ of all weak limits of sequences $\mu(dz, f_j)$, $f_j : U \rightarrow D$, $|f_j| < M$, and $\lim f_j(0) = 0$, is convex. Let $\mu(dz, f_j)$

converges to μ and $\mu(dz, g_j)$ converges to ν . We shall prove that $\alpha\mu + (1 - \alpha)\nu \in \mathbf{P}$. We may assume that all mappings f_j and g_j are holomorphic in a neighbourhood of $\overline{U}$ and $f_j(0) = g_j(0) = 0$. We consider mappings

$$h_j(\zeta) = f_j(\zeta) + g_j\left(\frac{r_j}{\zeta}\right),$$

defined on rings $R_j = \{\zeta : r_j \leq \zeta \leq 1\}$. Since either $|\zeta|$ or $r_j/|\zeta|$ is less than $\sqrt{r_j}$ on R_j , we see that for every j and every $\varepsilon_j > 0$ the set $h(R_j)$ belongs to the ε_j -neighborhood of $f_j(U) \cup g_j(U)$ when r_j is sufficiently small. In particular, we can find r_j such that $h_j(R_j) \subset D$. Let $e_j(\zeta)$ be the composition of e^ξ and conformal equivalence of the unit disk and the strip $\{\xi : \ln r_j < \mathbf{Re} \xi < 0\}$, mapping the origin into $\xi = (1 - \alpha) \ln r_j$ and the point $\zeta = 1$ into the origin. Evidently, mappings e_j map the arc $\gamma_1 = \{e^{i\theta} : -\pi\alpha < \theta < \pi\alpha\}$ onto the unit circle S and its complement γ_2 on the circle $r_j S$. We consider mappings $p_j = h_j \circ e_j$ of U into D .

If φ is a continuous function on $\mathbb{C}^n$ then

$$\int \varphi(z) \mu(dz, p_j) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(h_j(e_j(e^{i\theta})) + g_j(r_j e_j^{-1}(e^{i\theta}))) d\theta.$$

Because of symmetry it is enough to estimate the integral

$$\frac{1}{2\pi} \int_{-\pi\alpha}^{\pi\alpha} \varphi(f_j(e_j(e^{i\theta})) + g_j(r_j e_j^{-1}(e^{i\theta}))) d\theta.$$

Since $|g_j(r_j e_j^{-1}(e^{i\theta}))| \leq M r_j$ for $\theta \in \gamma_1$ the previous integral is equal to

$$\frac{1}{2\pi} \int_{-\pi\alpha}^{\pi\alpha} \varphi(f_j(e_j(e^{i\theta}))) d\theta + \delta_j,$$

where the sequence of numbers $\{\delta_j\}$ converges to 0 as $j \rightarrow \infty$. If u_j are harmonic functions on the unit disk with boundary values $\varphi(f_j(e^{i\theta}))$ then they are uniformly bounded and, therefore, $|u_j(\zeta) - u_j(0)| < K|\zeta|$. Thus,

$$u_j(e_j(0)) = \frac{1}{2\pi} \int_0^{2\pi} u_j(e_j(e^{i\theta})) d\theta = \frac{1}{2\pi} \int_{-\pi\alpha}^{\pi\alpha} \varphi(f_j(e_j(e^{i\theta}))) d\theta + \frac{1}{2\pi} \int_{\gamma_2} u_j(e_j(e^{i\theta})) d\theta.$$

Since $|u_j(e_j(0)) - u_j(0)| < Kr_j^{1-\alpha}$ and $|u_j(e_j(e^{i\theta})) - u_j(0)| < Kr_j$ for $\theta \in \gamma_2$, we see that

$$\frac{1}{2\pi} \int_{-\pi\alpha}^{\pi\alpha} \varphi(f_j(e_j(e^{i\theta}))) d\theta = \alpha u_j(0) + k_j,$$

where numbers k_j converge to 0. Therefore,

$$\int \varphi(z) \mu(dz, p_j) \rightarrow \alpha \int \varphi(z) \mu_1(dz) + (1 - \alpha) \int \varphi(z) \mu_2(dz),$$

and we proved the convexity of $\mathbf{P}$.

Clearly, the set $\mathbf{P}$ is closed. If μ is a Jensen measure with barycenter $0 \in D$ which is not in $\mathbf{P}$ then there is a continuous function φ on $\mathbb{C}^n$ such that

$$\int \varphi(z) \mu(dz) = a < b \leq \int \varphi(z) \nu(dz)$$

for all measures $\nu \in \mathbf{P}$. It was proved in [8] (see, also, [3]) that if $v(z)$ is a maximal plurisubharmonic function on D which is less than φ then the infimum over all $\nu \in \mathbf{P}$ of

$$\int \varphi(z) \nu(dz)$$

is equal to $v(0)$. Since any plurisubharmonic function can be approximated by continuous plurisubharmonic functions on compacts and since the support of μ is a compact in D we see that

$$a = \int \varphi(z) \mu(dz) \geq \int v(z) \mu(dz) \geq v(0) \geq b,$$

and we get the contradiction. $\square$

§3 LEAVES

Let $L = \{f_j\}$ be an uniformly bounded sequence of holomorphic mappings of the unit disk into $\mathbb{C}^n$, i.e. all sets $f_j(\overline{U})$ belong to a ball $D = B(0, R)$ for some $R > 0$. We introduce the cluster $\text{cl } L$ of L , as the set of all points $z \in \mathbb{C}^n$ such that there is a sequence of points $\zeta_{j_k} \in U$ with $\lim_{k \rightarrow \infty} f_{j_k}(\zeta_{j_k}) = z$.

Clusters can be quite messy and pathological. Our goal is to find inside of them something nice, and to make the first move at this direction we shall use weak compactness of holomorphic measures.

By the Alouglou theorem [5, V.4.2], the set of measures $\mu(dz, f_j)$ is compact with respect to the weak topology defined by integrals of continuous functions. Therefore, every sequence $L = \{f_j\}$ of uniformly bounded holomorphic mappings contains a subsequence $\{f_{j_k}\}$ with weakly converging holomorphic measures $\mu(dz, f_{j_k})$. We limit ourselves to studies of such sequences.

A sequence of uniformly bounded holomorphic mappings $L = \{f_j\}$ is called a *leaf* if measures $\mu(dz, f_j)$ are weakly converging to the Jensen measure $\mu(dz)$ of L , which will be denoted by $\mu(dz, L)$. Evidently, centers z_{f_j} of analytic disks f_j converge to a point z_L , which we shall call *the center* of L .

So we see that through every point of a general cluster passes a leaf made from the same sequence. But leaves can still be quite pathological. For example, let h_j be a conformal mapping of U onto the strip $D_j = \{\zeta = x + iy \in \mathbb{C} : |x| < 1, |y| < 1/j\}$ and g_j be any sequence of holomorphic mappings of D_j (which can approximate any continuous function on $[-1, 1]$). Then the sequence $f_j = g_j \circ h_j$ has as its cluster any cluster of a sequence of continuous mappings of $[-1, 1]$, which can be extremely unpleasant. To avoid the consideration of such pathologies (which is not a great loss as we show later) we should introduce a couple of definitions.

A point $z \in \text{cl } L$ is called *nonessential* if there is a number $r > 0$ such that

$$\lim_{j \rightarrow \infty} \omega(0, f_j^{-1}(B^n(z, r)), U) = 0.$$

Other points in $\text{cl } L$ are called *essential*. We shall denote the set of essential points of L by $\text{ess } L$. Evidently, this set is closed.

Any leaf $L = \{f_j\}$ determines the set $\text{supp } L$ equal to the support of the measure $\mu(dz, L)$. Evidently $\text{supp } L$ is compact, every point $z \in \text{supp } L$ is essential, and $\text{supp } L = \{z_L\}$ if and only if $\text{ess } L = \{z_L\}$.

The following theorem claims that we can get rid of non-essential points without losing too much.

Theorem 3.1. *For every uniformly bounded sequence $L = \{f_j\}$ of holomorphic mappings of U into $\mathbb{C}^n$ there is another sequence of such mappings $M = \{g_j\}$ such that:*

- (1) $\text{cl } M$ is the set of all essential points of $\text{cl } L$;
- (2) all points of $\text{cl } M$ are essential;
- (3) if L is a leaf then M is a leaf and $\mu(dz, L) = \mu(dz, M)$.

Proof. For every nonessential point z of $\text{cl } L$ we can find a ball $B^n(z, r(z))$ such that

$$\lim_{j \rightarrow \infty} \omega(0, f_j^{-1}(B^n(z, 2r(z))), U) = 0.$$

Let us choose a countable covering of the set of nonessential points by balls $B_k = B(z_k, r_k)$, where z_k is a nonessential point and $r_k = r(z_k)$. We denote by C_m the union of first m closed balls $\overline{B}_k$, $k = 1, \dots, m$. For every integer m there is an integer $j(m)$ such that

$$\omega(0, f_j^{-1}(C_m), U) < \frac{1}{m}$$

when $j \geq j(m)$. For $j(m) \leq j < j(m+1)$ we consider open sets $D_{mj} = f_j^{-1}(\mathbb{C}^n \setminus C_m)$ in U . Each of D_{mj} contains the origin and let p_j be an universal holomorphic covering mapping of D_{mj} by U such that $p_j(0) = 0$.

The mapping p_j has radial limits a.e. on S and, by Lindelöf theorem, $g_j(\zeta) = f_j(p_j(\zeta))$ for almost all points $\zeta \in S$. If E is a subset of U then, evidently,

$$(1) \quad \omega(0, p_j^{-1}(E), U) \geq \omega(0, E, U) - \frac{1}{m}.$$

We define $g_j = f_j \circ p_j$ and let $M = \{g_j\}$. Evidently, $\text{cl } M$ is the set of all essential points of L .

If z is an essential point of L then for every ball $B(z, r)$, $r > 0$ there is a subsequence $\{f_{j_k}\}$ such that

$$\omega(0, f_{j_k}^{-1}(B(z, r)), U) > a(r) > 0.$$

By (1)

$$\omega(0, g_{j_k}^{-1}(B(z, r)), U) > a(r) - \frac{1}{m} \rightarrow a(r) > 0.$$

So all points of $\text{cl } M$ are essential. If L is a leaf, the same reasoning shows that $\mu(dz, g_j)$ weakly converge to the same measure $\mu(dz, L)$ and the sequence $M = \{g_j\}$ is a leaf with $\mu(dz, M) = \mu(dz, L)$. $\square$

As the first application this theorem tells us that the set of essential points is quite big.

Corollary 3.1. *For every leaf L the set $\text{ess } L$ is connected.*

Proof. By Theorem 3.1 $\text{ess } L = \text{cl } M$ which is connected. $\square$

A sequence $M = \{g_j\}$ is called *perfect* if all points of $\text{cl } M$ are essential. A perfect leaf M satisfying conclusions of Theorem 3.1 for some sequence L is called a *perfection* of L .

Let V be an open set in $\mathbb{C}^n$ and $L = \{f_j\}$ be a sequence. For a point $z \in f_j(U)$ we define the function $\omega(z, V, f_j(U))$ to be equal to the maximum of $\omega(\zeta, f_j^{-1}(V), U)$ for all ζ such that $f_j(\zeta) = z$. We set $\omega(z, V, f_j(U))$ to be equal to 0 if $z \notin f_j(U)$. We introduce the harmonic measure on clusters as an upper semicontinuous function

$$\omega(z, V, L) = \overline{\lim}_{w \rightarrow z, j \rightarrow \infty} \omega(w, V, f_j(U))$$

on $\text{cl } L$.

If E is arbitrary set in $\mathbb{C}^n$ we let

$$\omega(z, E, L) = \inf \omega(z, V, L),$$

where the infimum is taken over all open sets V containing E .

Lemma 3.1. *If a point $z \in \text{cl } L$ is essential then for every open set $E \subset \text{cl } L$, containing z , $\omega(z_L, E, L) > 0$.*

Proof. Let E be an open set in $\text{cl } L$. We can find $r > 0$ such that the intersection of the ball $B_2 = B(z, 2r)$ and $\text{cl } L$ belongs to E . For an arbitrary $\delta > 0$ there is an open set $V \subset \mathbb{C}^n$ such that $V \cap \text{cl } L = E$ and $\omega(z_L, V, L) < \omega(z_L, E, L) + \delta$. There is an integer k such that $f_j(\zeta) \in V$ if $f_j(\zeta) \in B(z, r)$ and $j > k$. Therefore, $\omega(z_L, V, L) > \omega(z_L, B(z, r), L)$. Since z is essential, $\omega(z_L, B(z, r), L) > 0$. $\square$

Theorem 3.2. *Let $L = \{f_j\}$ be a perfect leaf and $X = \text{supp } L$. Then the harmonic measure $\omega(z, X, L) \equiv 1$ on $\text{cl } L$.*

Proof. Let V be an open neighbourhood of X . We must prove that for every $z \in \text{cl } L$ and for every neighbourhood W of z there is a sequence of points $a_{j_k} \in U$ such that:

- (1) $f_{j_k}(a_{j_k}) \in W$;
- (2)

$$\lim_{k \rightarrow \infty} \omega(a_{j_k}, f_{j_k}^{-1}(V), U) = 1.$$

This is evident if $z_L \in W$ because $\omega(z_L, V, L) = \mu(\mathbb{C}^n, L) = 1$. So we may assume that $z_L \notin W$.

Let $W_j = f_j^{-1}(W)$ and let $B_1 \Subset W$ be a ball centered in z . Since z is essential there is a subsequence j_k such that for sets $Z_k = f_{j_k}^{-1}(\overline{B_1})$ the harmonic measure

$$\omega(0, Z_k, U) > \delta > 0.$$

Let $V_k = f_{j_k}^{-1}(V) \cap S$ and $v_k(\zeta) = \omega(\zeta, V_k, U)$. Suppose that v_k is less than $\varepsilon < 1$ on W_{j_k} . If g_k is an universal cover of $U \setminus Z_k$ such that $g_k(0) = 0$, then radial limits of g_k are in Z_k on a set of measure greater than δ . Therefore, for the function $u_k = v_k \circ g_k$ we have

$$u_k(0) < (1 - \delta) + \varepsilon\delta = 1 - \delta(1 - \varepsilon) < 1.$$

But the sequence $u_k(0) = v_k(0)$ converges to 1, so there are an integer k and a point $a_{j_k} \in W_{j_k}$ such that

$$\omega(a_{j_k}, f_{j_k}^{-1}(V), U) > \varepsilon.$$

Since $\varepsilon < 1$ is arbitrary we get our theorem. $\square$

A point $z \in \text{cl } L$ is called *totally essential* if for every ball $B_1 = B(z, r)$ the lower limit of the sequence $\omega(0, f_j^{-1}(B_1), U)$ is greater than 0. Evidently, the set of totally essential points is closed and all points in $\text{supp } L$ are totally essential. A leaf is *totally perfect* if all point of its cluster are totally essential.

Theorem 3.3. *If $L = \{f_j\}$ is a perfect leaf then for every $z \in \text{cl } L$ there is a totally perfect leaf M such that $z_M = z_L$, $z \in \text{cl } L$, $\mu(dz, M) = \mu(dz, L)$, and $\text{cl } M$ belongs to $\text{cl } L$ and contains all totally essential points of L .*

Proof. Since z is an essential point of L , it is easy to see that there is a subsequence $\{f_{j_k}\}$ such that z is totally essential for this subsequence. Its perfection contains z and the set T of all totally essential points of L , which will continue to be totally essential. So we may assume that z is a totally essential point of L .

Let $n_1 > 1$ be the minimal positive integer such that the set

$$E_1 = \{z \in \text{cl } L : \text{dist}(z, T) \geq \frac{1}{n_1}\}$$

is non-empty. Since E_1 is compact there is a finite cover of E_1 by balls B_{1m} of radius less $1/2n_1$, centered at points of E_1 and such that for each ball B_{1m} there is a subsequence

$\{f_{j_k}\}$ with

$$\lim_{k \rightarrow \infty} \omega(0, f_{j_k}^{-1}(B_{1m}), U) = 0.$$

We take such a subsequence $\{f_{1j}\}$ for the ball B_{11} . For its perfection L_1 the set T_1 of all totally essential points contains T and the $\text{cl } L_1$ does not intersect B_{11} . If the ball B_{12} contains points of T_1 we leave it at peace, otherwise we repeat the procedure for the sequence $\{f_{1j}\}$. After finite number of steps we exhaust all balls B_{1m} and come to the leaf L_l with the set T_l of totally essential points. Every point of $\text{cl } L_l$ lies within the distance $1/n_1$ from points of T_l . So if n_2 is the minimal positive integer such that the set

$$E_2 = \{z \in \text{cl } L_l : \text{dist}(z, T_l) > \frac{1}{n_2}\}$$

is non-empty, then $n_2 > n_1$. We again take a finite cover by balls of radius less than $1/2n_2$ and repeat the previous construction.

At the end of the process we get a sequence of leaves $M_k = \{f_{kj}\}$ with perfections L_k and the sequence of balls B_k such that M_1 is a subsequence of L , M_{k+1} is a subsequence M_k , the ball B_k is non-essential for M_k . Sets T_k are increasing and their union is dense in the intersection of clusters of L_k . We take the sequence $\{g_j = f_{jj}\}$ and let M be its perfection.

First of all, $z_M = z_L$ and $\mu(dz, M) = \mu(dz, L)$. Secondly, $\text{cl } M$ belongs to the intersection of clusters of L_k and, therefore, doesn't contain the union of balls B_k . And if the point $z \in T_k$ for some k then it is totally essential for $\{g_j\}$. Therefore, $\text{cl } M$ consists from totally essential points only. $\square$

Let us give an example of a perfect but not a totally perfect leaf.

Example 3.1. Let $f_j = (\zeta, \zeta^j)$ be a sequence of mappings of U into U^2 . It is easy to see that measures $\mu(dz, f_j)$ converge to the measure

$$\mu = \frac{1}{2\pi} d\alpha d\beta$$

on $S \times S$. Therefore, the sequence of f_j determines the leaf $L = \{f_j\}$ with $\mu(dz, L) = \mu$. So $\text{supp } L = S \times S$, and

$$\text{cl } L = \{(z_1, z_2) : |z_1| < 1, z_2 = 0 \text{ or } |z_1| = 1, |z_2| \leq 1\}.$$

The leaf L is totally perfect.

We can consider another sequence of holomorphic mappings g_j equal to f_j if j is even, and to (ζ^j, ζ) if j is odd. Then holomorphic measures $\mu(dz, g_j)$ also converge to μ , so the sequence $M = \{g_j\}$ is a leaf, and this leaf is perfect but not totally perfect. Its cluster is the union of two perfect leaves.

§4. GEOMETRY AND ANALYTIC STRUCTURES ON LEAVES

If $L = \{f_j\}$ is a leaf in $\mathbb{C}^n$ then it is interesting to find out what does $\text{cl } L$ remember about its holomorphic origin. There is no sense to look at leaves with trivial Jensen measures: their clusters can be as pathological as clusters of continuous mappings. So we assume that $\text{supp } L \neq \{z_L\}$. In this case, the set $\text{ess } L \neq \{z_L\}$ and, by Theorem 3.1, we can find a perfect leaf with the same support and the same Jensen measure, which cluster is $\text{ess } L$. Since remaining points again can form extremely pathological set we shall concentrate on studies of perfect leaves.

The first question which naturally arises is the question about the existence of analytic disks in clusters. The negative answer to this question was given by Stolzenberg in [12] and Wermer in [14]. There are perfect leaves such that any holomorphic mapping of the unit disk into the cluster of such leaves is constant. Before we go further, we give the brief description of these examples.

Examples of Stolzenberg and Wermer. Both examples deal with complex analytic 1-dimensional irreducible varieties V_j (Stolzenberg) and W_j (Wermer) in the unit bidisk in $\mathbb{C}^2$. Each of V_j or W_j contains the origin. So, according to our approach, these sequences of varieties or their subsequence are leaves with supports lying on the boundary of U^2 . It was proved by Stolzenberg and Wermer that clusters of these leaves does not contain an analytic disk.

There is a difference in this examples. The example of Stolzenberg is a "Swiss cheese" lifted to $\mathbb{C}^2$, the projections of its cluster on both coordinate planes are nowhere dense. But if one makes a "Swiss cheese" on a plane then it may happen that its cluster (what is left after we drag out holes) consists from non-essential points only. By Theorem 3.1 Stolzenberg's example has a connected set of essential points joining the origin with the boundary of U^2 .

Varieties W_j in Wermer's example have proper projections on one of coordinate lines, say, $\{w = 0\}$. Moreover, there is a countable dense set of points $\{p_k\}$ on this line such that for each j the variety W_j has finite number of points over each of p_k , and there

is only one point in W_j over the origin. This property was used by Goldmann in [6] to show that the cluster of Wermer's example is perfect. In the same paper Goldmann discovered that this cluster has some analytic properties, which we discuss later. This discovery gave us the starting push for our studies. But before we move to this questions we want to give a sufficient conditions for the existence of analytic disks in clusters.

Theorem 4.1. *Let $L = \{f_j\}$ be a leaf in $\mathbb{C}^n$. Suppose that there is a holomorphic function h , defined on a neighbourhood V of $\text{cl } V$, and numbers $b < 1$ and $\alpha > 0$ such that:*

- (1) $|h|$ and $\|\nabla h\|$ are less than $M < \infty$ on V ;
- (2) $|h(z_L)| \leq b < 1$;
- (3) for every integer $j \geq 1$ there is an arc γ_j with the length greater than α such that

$$\lim_{r \rightarrow 1} |h(f_j(r\zeta))| \geq 1$$

for all points ζ in γ_j .

Then there is a non-constant holomorphic mapping $g = (g_1, \dots, g_n) : U \rightarrow \text{cl } L$ such that at least one of functions g_k , $1 \leq k \leq n$, covers univalently a disk of a positive radius $r = r(M, b, \alpha)$ in $\mathbb{C}$.

Proof. Let $S(c)$ be a set of holomorphic functions u on U with the Bloch norm

$$\|u\|_B = \sup_{\zeta \in U} |u'(\zeta)|(1 - |\zeta|^2) \geq c.$$

By Theorems 2 and 1 from [7] functions $h_j = h \circ f_j$ belong to $S(c)$ for some $c = c(M, b, \alpha) > 0$. Since $\|\nabla h\| < M$ there is an integer $k(j)$ such that

$$\|f_{jk(j)}\|_B \geq \frac{c}{nM},$$

where f_{jk} is the k -th coordinate function of f_j . So for some k between 1 and n we can find an infinite subsequence f_{j_i} such that:

(1)

$$|f'_{j_i k}|(1 - |\zeta_{j_i}|^2) \geq \frac{c}{2nM}$$

at some point ζ_{j_i} in U ;

(2) the sequence $f_{j_i}(\zeta_{j_i})$ converges to z_0 .

Mappings

$$f_{j_l} \left(\frac{\zeta - \zeta_{j_l}}{1 - \bar{\zeta}_{j_l} \zeta} \right)$$

converge to a non-constant mapping g which image lies in $\text{cl } U$ and $|g'_k(0)| \geq c/2nM$. So, by the Bloch theorem, g_k covers univalently a disk of a positive radius $r = r(M, b, \alpha)$. $\square$

Examples of Stolzenberg and Wermer (see, also, [7]) show that it is hard to await something really good at this situation but in the one-dimensional case we can do a little bit better.

Theorem 4.2. *Let $L = \{f_j\}$ be a perfect leaf in $\mathbb{C}$ and Δ be the boundary of $\text{cl } L$. Then $\Delta \subset \text{supp } L$.*

Proof. Suppose that there is a disk $B_1 = B(z_0, r)$ such that $z_0 \in \Delta$, $r > 0$ and $\mu(B_1, L) = 0$. We can find a point z such that a closed disk $\bar{B}_2 = \bar{B}(z, s) \subset B(z_0, r/4)$ does not intersect $\text{cl } L$. Therefore, there is an integer N such that $f_j(U) \cap B_2 = \emptyset$ when $j > N$. Let $B_3 = B(z, r/2)$ and let $u(z)$ be a harmonic function on $\mathbb{C} \setminus B(z, s)$, equal to 1 on ∂B_2 and to 0 on ∂B_3 . Functions $u_j(\zeta) = u(f_j(\zeta))$ will be harmonic on U . If $E_j = \{\zeta \in S : u_j(\zeta) > 0\}$ then the sequence of $a_j = m(E_j)$ tends to 0. If $B_3 = B(z_0, r/4)$ then $u(z) > c > 0$ on B_3 . Harmonic functions v_j on U with boundary values equal to 1 on E_j and to 0 on $S \setminus E_j$ are greater than u_j . The harmonic measure of the set $F_j = \{\zeta \in U : v_j(\zeta) > c\}$ is less than a_j/c and, therefore, the harmonic measure of sets $G_j = \{\zeta \in U : u_j(\zeta) > c\}$ is less than a_j/c which means that the point z_0 is nonessential. $\square$

In general, it may happen that $\mu(\Delta \cap B(z, r), L) = 0$ for some $z \in \Delta$.

Corollary 4.1. *Let $L = \{f_j\}$ be a leaf in $\mathbb{C}$ such that $z_L = 0$ and $\text{supp } L$ lies outside the disk $B_1 = B(0, r)$ for some $r > 0$. Then B_1 belongs to $\text{cl } L$.*

Proof. We may assume that the leaf L is perfect. Since $\text{supp } L$ does not intersect B_1 , the boundary of $\text{cl } L$ does not intersect B_1 . So B_1 belongs to $\text{cl } L$. $\square$

Example 4.1. Let $f_j = (g_j, h_j)$ be uniformizations of curves V_j from the Stolzenberg's example by analytic disks such that $f_j(0) = (0, 0)$. We may assume that $\{f_j\}$ is a leaf L . Then $\text{supp } L \subset \partial U^2$ and, therefore, either the leaf $G = \{g_j\}$ or the leaf $H = \{h_j\}$ is non-trivial. Their perfections are also non-trivial. Nevertheless, their clusters do not contain any disk in $\mathbb{C}$. In this example, $\text{cl } L = \text{supp } L = \Delta$.

So the complex geometry is not inherited by leaves explicitly. To go further we introduce for each leaf L algebras $\mathcal{P}(L)$ and $\mathcal{H}(L)$ of functions on $\text{cl } L$, which can be approximated on $\text{cl } L$ either by polynomials or by functions holomorphic in a neighbourhood of $\text{cl } L$. Such algebras are called *analytic* if a function from the algebra is equal to 0 on $\text{cl } L$ provided it is equal to 0 on some open set V in $\text{cl } L$.

Our first step is to prove the two constant theorem for functions in $\mathcal{H}(L)$.

Theorem 4.3. *Let $L = \{f_j\}$ be a sequence of uniformly bounded analytic disks and let $h \in \mathcal{H}(L)$. If $|h|$ is less than M on $\text{cl } L$ and less than $m < M$ on an a set $E \in \text{cl } L$, then*

$$|h(z)| < m^d M^{1-d},$$

where $d = \omega(z, E, L)$.

Proof. Let us fix some $\varepsilon > 0$. The function h is the limit of functions h_k holomorphic in neighbourhoods D_k of $\text{cl } L$. We may find an integer k such that $|h_k| < M' = (1 + \varepsilon)M$ on D_k , $|h_k| < m' = (1 + \varepsilon)m$ on E , and $|h_k(z) - h(z)| < \varepsilon$.

Let V be the open set of points in D_k , where $|h_k|$ is less than m' . Evidently, $\omega(z, V, L) \geq d$. If $V_j = f_j^{-1}(V)$ then, by the definition of $\omega(z, V, L)$, there is a sequence of points $a_{j_l} \in U$ such that points $f_{j_l}(a_{j_l})$ converge to z and the limit d' of the sequence $d_{j_l} = \omega(a_{j_l}, V_{j_l}, U)$ is greater than d . We may assume that $j_l = j$.

If $v_j = h_k \circ f_j$ then by the two constants theorem

$$|v_j(a_j)| < (m')^{d_j} (M')^{1-d_j}$$

and

$$|h_k(z)| < (m')^{d'} (M')^{1-d'} \leq (m')^d (M')^{1-d}.$$

Therefore, $|h(z)| < m^d M^{1-d}$. $\square$

Corollary 4.2. *Let $L = \{f_j\}$ be a perfect leaf in $\mathbb{C}^n$ and let z_L be the center of L . If a function $h \in \mathcal{H}(L)$ is equal to zero on an open set $E \subset \text{cl } L$ then $h(z_L) = 0$.*

Proof. This statement follows from the previous theorem because all points in $\text{cl } L$ are essential and, therefore, $\omega(z_l, E, L) > 0$. $\square$

Corollary 4.3. *Let $L = \{f_j\}$ be a perfect leaf in $\mathbb{C}^n$. Then:*

- (1) *every function $h \in \mathcal{H}(L)$ attains its maximum modulus value on $\text{supp } L$;*
- (2) *cluster of L belongs to the polynomial hull of $\text{supp } L$.*

Proof. Part 1) follows immediately from Theorem 4.3 and 3.2 and part 2) is the consequence of part 1). $\square$

The maximum modulus principle proved just now can be obtained from the fact that the measure $\mu(dz, L)$ is an Jensen measure for $\mathcal{H}(L)$ at z_L . We did not explore this approach because we intended to be as geometric as possible. Connections of this subject with the theory of uniform algebras are evident. Nevertheless, it seems to us that this theory does not allow to look closely at geometric structures generated by leaves. Moreover, we shall prove in §5 that polynomial hulls can be foliated by perfect leaves, so our method can be applied to uniform algebras as well. Let us consider the following example.

Example 4.2. Let L be a perfect leaf from Example 3.1. The function $h(z_1, z_2) = w$ is equal to 0 on a neighbourhood of origin and is not 0 everywhere. So perfect leaves lack of the propagation of zeros. To understand this phenomenon we must look closer at the geometry of leaves.

It happens that leaves generate complex substructures in a natural way which we are going to describe now. Let

$$G_a(\zeta) = \frac{\zeta + a}{1 + \bar{a}\zeta}$$

be a conformal automorphism of the unit disk. Let $L = \{f_j\}$ be a perfect leaf. For a point $z \in \text{cl } L$ we consider points $a_k \in U$ such that the a subsequence $f_{j_k}(a_k)$ converges to z . From the sequence of mappings $g_k = f_{j_k} \circ G_{a_k}$ we can choose a subsequence h_m with weakly converging measures $\mu(dz, h_m)$. A *subleaf* L_z is the perfection of the leaf $M = \{h_k\}$.

If M is a subleaf of L then $\text{cl } M \subset \text{cl } L$ but it may happen that $\text{supp } M$ is not a subset of $\text{supp } L$. For example, if f_j are an universal holomorphic coverings of $U \setminus \overline{B}(1/2, 1/j)$ by U such that $f_j(0) = 0$, then there are subleaves $M = L_{\frac{1}{2}}$ such that $\mu(\frac{1}{2}, M) = a$, where a is any number between 0 and 1. A subleaf L_z such that $\text{supp } L_z \subset \text{supp } L$ is called a *principal subleaf*.

The set of all subleaves of the leaf L is partially ordered by the inclusion relation of their clusters. A leaf L is called *minimal at a point* $z \in \text{cl } L$ if it contains no proper non-trivial subleaves passing through z . Evidently, a leaf L , minimal at z_L , is totally perfect. A leaf L is called *minimal* if it is minimal at all points of $\text{cl } L$. A point $z \in \text{cl } L$ is called *inner* if there is a non-trivial subleaf L_z . We shall say that a sequence of leaves

$\{L_j\}$ converges to a leaf L if every neighbourhood of every point of $\text{cl } L$ contains points from clusters of infinitely many different leaves L_j and $\mu(dz, L_z) = \lim \mu(dz, L_j)$

In Example 4.2 there is only one subleaf passing through the origin, so L is minimal at z_L . The leaf M from Example 3.1 has two subleaves passing through the origin so it is not minimal at z_L . Both leaves have subleaves at the boundary of the bidisk so they are not minimal. These subleaves are analytic disks foliating the boundary. If we take the leaf L and a point $z_0 = (z_1, z_2)$ such that $|z_1| = 1$ and $z_2 \neq 0, |z_2| < 1$, then for each j we can find a point $a_j \in U$ such that $f_j(a_j)$ converges to z_0 and let $g_j = f_j \circ G_{a_j}$. We may assume that the sequence g_j is a leaf $L' = \{g_j\}$, otherwise we take a converging subsequence. Evidently, $\text{cl } L = \text{cl } L'$, but the leaf L' is not perfect. The origin is a non-essential point of M . We shall show that the property for a leaf to be minimal and the analyticity of the algebra $\mathcal{H}(L)$ are closely related.

Lemma 4.1.

- (1) *If points $z_j \in \text{cl } L$ converge to a point z and measures $\mu(dz, L_{z_j})$ weakly converge to a measure μ then subleaves L_{z_j} converge to a subleaf L_z ;*
- (2) *if J is a totally ordered set and $\mathcal{L} = \{L_{z_j}, j \in J\}$ is a set of subleaves such that $\text{cl } L_{z_i} \subset \text{cl } L_{z_j}$ for $i > j$, then there is a countable sequence $\{L_j\}$ in $\mathcal{L}$ converging to a subleaf which belongs to all subleaves in $\mathcal{L}$;*
- (3) *if z belongs to the cluster of a perfect leaf L but doesn't belong to its support then there is a principal subleaf L_z passing through z .*

Proof. 1) If leaves $L_{z_j} = \{g_{jk} = f_j \circ G_{a_{jk}}\}$ then we can find a subsequence $a_m = a_{j_m k_m}$ in U such that measures $\mu(dz, g_{j_m k_m})$ converge to the measure μ . Then the sequence $f_{j_m} \circ G_{a_m}$ defines a leaf L_z .

2) Let G be the intersection of clusters of all leaves in $\mathcal{L}$. There is a countable sequence j_k such that the set G is the intersection of $\text{cl } L_{j_k}$. Let us select another subsequence with converging holomorphic measures. This subsequence of subleaves converges to a subleaf L_z due to the first part of the lemma.

3) Suppose that there is a point $z \in \text{cl } L$ such that $\text{dist}(z, \text{supp } L) = r > 0$ and for every L_z the support of L_z doesn't belong to $\text{supp } L$. This means that there are numbers $a > 0$ and $d > 0$, and a ball B_1 , centered at z , such that

$$\lim_{j \rightarrow \infty} \mu(W, f_j \circ G_\zeta) > a$$

for every point w in B_1 and every $\zeta \in f^{-1}(w)$ and for the set W of points in $\mathbb{C}^n$, with distances to $\text{supp } L$ greater or equal to d . Otherwise, we can find a subsequence $f_{j_k} = g_k$, a sequence $w_k \in \mathbb{C}^n$ converging to z , and a sequence $\zeta_k \in g_k^{-1}(w_k)$ such that for the complement V of $\text{supp } L$ measures $\mu(V, g_k \circ G_{\zeta_k})$ converge to 0, and, therefore, there is a subleaf with the support lying in the support of L .

Let E_j be a subset of S such that $f_j(\zeta) \in W$ for every $\zeta \in E_j$. If $D_j = f_j^{-1}(B_1)$ then the function $\omega(\zeta, E_j, U)$ is greater than a on D_j when j is big enough. Since z is an essential point of L then $\omega(0, D_j, U) > \varepsilon > 0$ for infinitely many j . Therefore, $\omega(0, E_j, U) > \varepsilon a$ for such j , which means that $\mu(W, L) > \varepsilon a > 0$. $\square$

Inner points on leaves play the role close to the role of inner points in domains. For example, if D is a domain in $\mathbb{C}$ and f is a mapping of U onto D then for the leaf $L = \{f\}$ the inner points are points in the interior of D . The third part of Lemma 4.1 also shows that every point of $\text{cl } L$, which does not belong to the $\text{supp } L$, is inner. In particular, for examples of Stolzenberg and Wermer all points of clusters lying in U^2 are inner.

Corollary 4.4. *If z is a point of a perfect leaf L which does not belong to $\text{supp } L$ then there is a minimal non-trivial leaf L_z .*

Proof. We consider the set of all principal subleaves passing through z . By Lemma 4.1 every totally ordered subset of this set has a minimal element. So, by Zorn lemma, there is a minimal element in this set. $\square$

Lemma 4.2. *Let $f : U \rightarrow \mathbb{C}^n$ be a holomorphic mapping such that $\|f\| < R$, $f(0) = z_0$, $b = \|z_0\| \neq 0$, and the harmonic measure $\omega(z_0, B_1, f(U)) = a > 0$ for a ball $B_1 = B(0, r)$, $r < b$. Then for every $k > 1$ such that $kr < b$ there are a number $c = c(k, r, R, a, b) > 0$, a number $m = m(k, r, R, a, b)$, $k > m > 1$, and a point $z_1 \in S(0, mr)$ where $\omega(z_1, B_1, f(U)) > c$ and $\omega(z_1, \mathbb{C}^n \setminus B(0, kr), f(U)) > c$.*

Proof. Let

$$u(z) = \frac{\ln \frac{\|z\|}{R}}{\ln \frac{r}{R}}.$$

The function u is plurisuperharmonic, positive on $B(0, R)$, and greater than 1 on B_1 . If $1 < m < k$ and $d = \omega(0, f^{-1}(B(0, mr), U))$ then

$$u(z_0) \geq d \frac{\ln \frac{mr}{R}}{\ln \frac{r}{R}}$$

or

$$d \leq \frac{\ln \frac{b}{R}}{\ln \frac{mr}{R}}.$$

Let $v(\zeta) = \omega(\zeta, f^{-1}(B_1), U)$ and let $\gamma = f^{-1}(S(0, mr))$. If q is the maximum of v on γ then $qd \geq a$ or

$$q \geq \frac{a}{d} = a \frac{\ln \frac{mr}{R}}{\ln \frac{b}{R}} \geq a \frac{\ln \frac{kr}{R}}{\ln \frac{b}{R}} = 2s.$$

So there is a point ζ_1 on γ where v is greater than s . Let $z_1 = f(\zeta_1)$.

Let $B_2 = B(0, kr)$ and $p = \omega(\zeta_1, f^{-1}(\mathbb{C}^n \setminus B_2), U)$. Then

$$u(z_1) = \frac{\ln \frac{mr}{R}}{\ln \frac{r}{R}} \geq s + (1 - s - p) \frac{\ln \frac{kr}{R}}{\ln \frac{r}{R}}$$

or

$$\ln \frac{mr}{R} \leq s \ln \frac{r}{R} + (1 - s) \ln \frac{kr}{R} - p \ln \frac{kr}{R} = (1 - s) \ln k + \ln \frac{r}{R} - p \ln \frac{kr}{R}.$$

Therefore,

$$p \geq \frac{(1 - s) \ln k - \ln m}{\ln \frac{kr}{R}}.$$

If

$$m = k^{(1-s)/2}$$

then

$$p \geq \frac{(1 - s) \ln k}{2 \ln \frac{R}{kr}} = t.$$

If we take $c = \min(s, t)$ we get the proof of the lemma. $\square$

Theorem 4.4. *For a perfect non-trivial leaf $L = \{f_j\}$:*

- (1) *the set of inner points of L is dense in $\text{cl } L$;*
- (2) *if a set V is open in $\text{cl } L$ and $z \in \text{cl } L$ then $\omega(z, W, L) = 0$ for every set $W \Subset V$ if and only if any non-trivial subleaf L_z does not intersect V .*

Proof. 1) We assume that $\|f_j\| < R/2$. Since L is non-trivial we can find a point $z_1 \in \text{cl } L$ such that $\|z_1 - z_L\| = b > 0$. For a ball $B_1 = B(z_1, r)$, $r > 0$, we consider the ball $B_2 = B(z_1, 2r)$. Since the leaf L is perfect we can find a subsequence f_{j_k} such that $\omega(z_L, B_1, f_{j_k}(U)) > a > 0$. By Lemma 4.2 there are points $\zeta_{j_k} \in U$ such that

$f_{j_k}(\zeta_{j_k}) = z_k \in S(z_1, mr)$, $1 < m < 2$, and $\omega(z_k, \mathbb{C}^n \setminus B_2, f_{j_k}(U)) > c > 0$. From the sequence

$$g_k = f_{j_k} \circ G_{\zeta_{j_k}}$$

we can select a subsequence forming a leaf $N = \{h_l\}$ and let M be the perfection of N . Then $z_M \in S(z_1, mr)$ and there are points of $\text{cl } M$ lying outside B_2 . Otherwise all these points are non-essential for the sequence h_l , so we can cover the compact intersection of $\mathbb{C}^n \setminus B_2$ and $\text{cl } N$ by a finite number of balls such that for each of them its harmonic measure at z_M tends to 0 as l goes to ∞ . This means that $\omega(z_M, \mathbb{C}^n \setminus B_2, f_{h_l}(U)) > c > 0$ tends to 0 and this is the contradiction. So the leaf M is non-trivial and, since r is arbitrary, we get the proof of the first part of the theorem. $\square$

2) If $\omega(z, W, L) = 0$ then for every sequence $g_k = f_{j_k} \circ G_{\zeta_{j_k}}$ such points $f_{j_k}(\zeta_{j_k})$ converge to z , all points in V are non-essential and, hence, cannot belong to the perfection of $\{g_k\}$. So every subleaf L_z does not intersect W and, therefore, V .

Conversely, if $\omega(z, W, L) > 0$ for some $W \subseteq V$ then there are essential points in $\overline{W}$ for the sequence $\{g_k\}$. Therefore, there are points of V in the perfection of this sequence. $\square$

Corollary 4.5. *If a perfect leaf L is minimal at z_L and z_L does not belong to $\text{supp } L$ then for every open set V in $\text{cl } L$ there is an open set W containing z_L such that every function $h \in \mathcal{H}(L)$ equal to 0 on V is equal to 0 on W .*

Proof. Suppose that there is a sequence of points z_k converging to z_L such that $h(z_k) \neq 0$. By Theorem 4.4 we may assume that points z_k are inner and, therefore, there are principal subleaves $L_k = L_{z_k} = \{f_{kj}\}$, which, by the second part of Theorem 4.4, don't intersect V . If we make the perfect leaf M from the sequence $\{f_{kj}\}$, then $z_M = z_L$ and M is non-trivial because $\text{supp } M$ belongs to $\text{supp } L$ and, hence, doesn't contain z_M . But $\text{cl } M$ is a proper subset because it doesn't contain V which contradicts to the minimality of L in z_L . $\square$

Theorem 4.5. *A leaf L is minimal at any point $z \in D$ if and only if $\omega(z, V, L) > 0$ for every inner point $z \in \text{cl } L$ and every open set $V \subset \text{cl } L$.*

Proof. If $\omega(z, V, L) = 0$ for some open V and some inner point z then, by Theorem 4.4, a non-trivial subleaf L_z doesn't intersect V and, hence, L is not minimal at z .

Conversely, if L is not minimal at z then there is an open set V which doesn't intersect some L_z . By the second part of Theorem 4.4 this implies that $\omega(z, V, L) = 0$. $\square$

Corollary 4.6. *If a leaf is minimal then the algebra $\mathcal{H}(L)$ is analytic.*

Proof. Follows immediately from Theorem 4.5, 4.4 1), and 4.3. $\square$

In his paper [6] Goldmann proves, in different terminology, that the leaf corresponding to the Wermer's example is minimal. So such leaves exist. Nevertheless, it seems that in general situation minimality does not happen frequently. If we consider the leaf L from Example 3.1 then the only part of this leaf where we can guarantee that the algebra $\mathcal{H}(L)$ is analytic is the disk $\{z_2 = 0\}$. To find an approach to this situation let us introduce the new definition. If $L = \{f_j\}$ is a leaf then *the midrib* of L is the closure of the set of all points z in $\text{cl } L$ such that $\omega(z, W, L) > 0$ for every neighbourhood W of z_L . For the leaf L mentioned above the midrib is the disk $\{z_2 = 0\}$.

Corollary 4.7. *If a perfect leaf L is minimal at z_L and z_L does not belongs to $\text{supp } L$ then every holomorphic function h on L equal to 0 on an open set V in $\text{cl } L$ is equal to 0 on the midrib of L .*

Proof. By Corollary 4.5 the function h is equal to 0 a neighbourhood W of z_L and, by two constants theorem, is equal to 0 on the midrib. $\square$

I do not know whether midribs are leaves or not, and I also do not know whether they are always nontrivial. The following theorem, which can be applied to the Wermer's example, gives sufficient criterion for the existence of a nontrivial midrib.

Theorem 4.6. *Let $L = \{f_j\}$ be a leaf and let h be a function from $\mathcal{H}(L)$ such that $h(z_L) \neq h(z)$ when $z \neq z_L$ is in $\text{cl } L$. We suppose that z_L does not belong to $\text{supp } L$. If K is a connected component, containing $h(z_L)$, of the complement of $\text{supp } L$ in $\mathbb{C}$ and $h(z) \in K$ for some point $z \in \text{cl } L$ then belongs to the midrib of L .*

Proof. We fix a neighbourhood W of z_L and a point $z \in \text{cl } L$ such that $h(z) \in K$. Let $W^* = h(W)$. Since $h(z_L) \neq h(w)$ for all $w \neq z_L$ we can find a neighbourhood $V \subseteq W$ of $h(z_L)$ such that $h^{-1}(V) \subseteq W$. Let us take an approximation g of h by functions holomorphic on $\text{cl } L$ such that $g(z)$ belongs to the connected component K' , containing $g(z_L)$, of the complement of the set $g(\text{supp } L)$ in $\mathbb{C}$ and $g^{-1}(V) \subseteq W$. We consider on K' the function

$$u(\zeta) = \omega(\zeta, g(W), K')$$

which is, evidently, greater than 0 at $g(z)$. Since

$$\omega(z, W, L) \geq u(g(z)) > 0$$

the point z belongs to the midrib. $\square$

It follows from this theorem that the leaf generated by the example of Wermer [14] coincides with its midrib.

§5. POLYNOMIAL HULLS

Let D be a domain in $\mathbb{C}^n$ and let ψ be an upper semicontinuous function on D . We consider the function

$$P\psi = u(z) = \inf \frac{1}{2\pi} \int_0^{2\pi} \psi(f(e^{i\theta})) d\theta,$$

where the infimum is taken over all mappings $f \in A(\overline{U}, D)$ with $f(0) = z$. It was proved in [8] that $u(z)$ is plurisubharmonic and

$$u(z) = \sup\{v(z) : v \leq \psi \text{ and } v \text{ is psh}\}.$$

Let $E \subset D$ be an open set. Then the characteristic function χ_E of E is lower semicontinuous and we define the *pluri-harmonic measure* $\omega(z, E, D)$ of the set E with respect to D to be equal to $P(-\chi_E)$. If F is an arbitrary set in D then we define the pluri-harmonic measure $\omega(z, F, D)$ to be equal to $\sup \omega(z, E, D)$, where the supremum is taken over all open sets E containing F . This function may be not upper semicontinuous, so we take its regularization

$$\omega^*(z, F, D) = \overline{\lim}_{w \rightarrow z} \omega(z, F, D),$$

which is plurisubharmonic.

If X is a compact in $\mathbb{C}^n$ then we shall denote by $\hat{X}$ the polynomial hull of X . It follows from Theorem 3.3 that if a leaf L is perfect and $X = \text{supp } L$ then $\text{cl } L \subset \hat{X}$.

If $X \subset D$ and D is a Runge domain then $\hat{K}$ coincides with the holomorphic envelope of K in D . The following theorem was proved in [9]. For the reader's convenience we shall supply the proof.

Theorem 5.1. *Let D be a Runge domain and K be a compact in D . Then $z_0 \in \hat{K}$ if and only if for every open set $E \supset K$ the pluri-harmonic measure $\omega(z_0, E, D) = -1$.*

Proof. Let $z_0 \in \hat{K}$ and E is an open set containing K . Then there is a decreasing sequence of continuous psh functions $v_k(z)$ on D converging to $\omega(z, E, D)$ pointwise on

D [11]. By [2] for every $\varepsilon > 0$ there are holomorphic functions $f_{jk}, j = 1, 2, \dots, N_k$ on D and positive numbers a_{jk} such that

$$\max_j \{a_{jk} \ln |f_{jk}|\} \leq v_k \leq \max_j \{a_{jk} \ln |f_{jk}|\} + \varepsilon$$

on K . Since D is a Runge domain we may assume that all f_{jk} are polynomials. We can take k so big that $v_k \leq -1 + \varepsilon$ on K . Then $v_k(z_0) \leq -1 + 2\varepsilon$ and this means that $\omega(z_0, E, D) = -1$.

Conversely, let g be a polynomial. For every $\varepsilon > 0$ we consider the open set $E = \{z \in D : |g(z)| < \sup_K |g(w)| + \varepsilon\} \supset K$. If $\omega(z_0, E, D) = -1$ then there is a mapping $f \in A(\overline{U}, D)$ with $f(0) = z_0$ such that $m\{\theta : f(e^{i\theta}) \in E\} > 2\pi(1 - \varepsilon)$. Therefore,

$$|g(z_0)| \leq \frac{1}{2\pi} \int_0^{2\pi} |g(f(e^{i\theta}))| d\theta \leq \sup_K |g(w)| + M\varepsilon,$$

where $M = \sup_D |g(w)|$. Hence, $z_0 \in \hat{K}$. $\square$

The following theorem establishes something like analytic structure in $\hat{X}$.

Theorem 5.2. *If X is a compact in $\mathbb{C}^n$ and $z \in \hat{X} \setminus X$ then there is a perfect leaf $L = \{f_j\}$ such that:*

- (1) $z = z_L$;
- (2) $\text{supp } L \subset X$;
- (3) $\text{cl } L \subset \hat{X}$;
- (4) *for every neighbourhood V of $\hat{X}$ there is an integer k such that analytic disks $f_j(U)$ are in V when $j > k$.*

Proof. By Theorem 5.1 we can find open neighbourhoods W_j of X with the intersection equal to X and analytic disks f_j such that the measure of points $\zeta \in S$ with $f_j(\zeta) \in W_j$ is greater than $2\pi - j^{-1}$ and $f_j(0) = z$. Taking, if necessary, a subsequence we may assume that the sequence $L = \{f_j\}$ is a leaf. Evidently, $\text{supp } L \subset X$. We may also assume that L is perfect, otherwise we replace L by its perfection.

By the construction, $z = z_L$ and $\text{supp } L \subset X$. Since $\text{cl } L$ belongs to the polynomial hull of $\text{supp } L$, we see that $\text{cl } L \subset \hat{X}$. Therefore, for every neighbourhood V of $\hat{X}$ there is an integer k such that analytic disks $f_j(U)$ are in V when $j > k$. $\square$

§6. MAXIMAL FUNCTIONS

The plurisubharmonic (psh) function u on a domain $D \subset \mathbb{C}^n$ is called *maximal* if for every domain $G \Subset D$ and every psh function v on D the inequality $v \leq u$ on ∂G implies the inequality $v \leq u$ on G . Bedford and Kalka proved in [1] that if a maximal function u is in $C^2(D)$ then through every point $z \in D$ we may draw a holomorphic disk such that the restriction of u to this disk is harmonic. The set of all this disks is called *the Monge–Ampère foliation for u* .

It was proved in [9, Lemma 2.2.3, Theorem 2.2.1] that maximal functions on strongly pseudoconvex domains can be obtained as solutions of variational problems for mappings of the unit disk. More precisely:

Theorem 6.1. *Let D be a strongly pseudoconvex domain in $\mathbb{C}^n$ and let $\varphi < M$ be a continuous function on ∂D . We let φ to be equal to M on D . Let*

$$\Phi(f) = \frac{1}{2\pi} \int_0^{2\pi} \varphi(f(e^{i\theta})) d\theta$$

be a functional on the space $A(U, D)$ of all holomorphic mappings $f : U \rightarrow D$. Then the function $u(z) = P\varphi = \inf \Phi(f)$, where the infimum is taken over all $f \in A(U, D)$ such that $f(0) = z$, is a maximal psh function on D , continuous up to the boundary, and equal φ on ∂D .

Moreover, for every point $z \in D$ there is a sequence $\{f_j\}$ of holomorphic mappings $f_j : U \rightarrow D$ such that $f_j(0) = z$, $\text{supp } \mu(dz, f_j)$ belongs to ∂D , and $\Phi(f_j) \rightarrow u(z)$ as $j \rightarrow \infty$.

As the following example shows the existence of Monge–Ampère foliations in the sense of Bedford and Kalka, is not valid in the general case.

Example 6.1. Let K be a compact in the boundary of the unit ball B , such that its polynomial hull $\hat{K} \subset B$ doesn't contain analytic disks. (One can use examples of Stolzenberg and Wermer to build such K .) If φ is a continuous negative function on $S = \partial B$ equal to -1 on K and greater than -1 everywhere else, then we can find a maximal function v which is continuous in $\overline{B}$ and is equal to φ on S . By Theorem 5.1 the function $v = P\varphi = -1$ only on $\hat{K}$. If $z \in \hat{K}$ and $f : U \rightarrow D$ is a holomorphic mapping such that $f(0) = z$ and $v \circ f$ is harmonic then $v \circ f \equiv -1$ and, hence, $f(U) \subset \hat{K}$. But

$\hat{K}$ does not contain analytic disks, so f is trivial, which means that the Monge–Ampère foliation for v does not cover $\hat{K}$.

So to extend the result of Bedford and Kalka to the general case we have to look for more general constructions.

Let $L = \{f_j\}$ be a leaf such that $\text{supp } L \subset \partial D$ and let $G \subset D$ be another domain, containing an inner point z of L . For a principal subleaf $L_z = \{g_j\}$ we consider a connected component G_j , containing the origin, of the set $g_j^{-1}(G)$ and let h_j be a holomorphic covering mapping of G_j by U such that $h_j(0) = 0$. Then the restriction of L on G at z is the perfection of the sequence $\{g_j \circ h_j\}$. We say that an upper semicontinuous function u on $\text{cl } L$ is *maximal* if for all domains G and H such that $G \Subset H \Subset D$ and every plurisubharmonic (resp. plurisuperharmonic) function v on H , the inequality $u \geq v$ (resp. $u \leq v$) on the intersection of $\text{cl } L$ and ∂G implies that $u(z) \geq v(z)$ (resp. $u(z) \leq v(z)$) on the intersection of $\text{cl } L$ and G .

Theorem 6.2. *Let D be a bounded domain in $\mathbb{C}^n$ and let u be a continuous plurisubharmonic function on D . We consider the functional*

$$\Phi(f) = \frac{1}{2\pi} \int_0^{2\pi} u(f(e^{i\theta})) d\theta$$

on the set $A(\overline{U}, D)$. Suppose that for a point $z_0 \in D$ there is a leaf $M = \{f_j\}$ such that $f_j \in A(U, D)$, $z_0 = z_M$, $\text{supp } M \subset \partial D$ and

$$u(z_0) = \lim_{j \rightarrow \infty} \Phi(f_j).$$

Then u is maximal on the perfection L of M .

Proof. We may assume that $L = M$. It is clear that $u(z) \leq \Phi(f)$ for every $z \in D$ and for every $f \in A(U, D)$ such that $f(0) = z$. Let us take arbitrary domains $G \Subset H \Subset D$ and let v be a plurisubharmonic function on H such that $u \geq v$ on the intersection K of $\text{cl } L$ and ∂G . We may assume that both u and v are less than some number A on H .

Suppose that there is a point z in the intersection F of $\text{cl } L$ and G such that $u(z) < v(z)$. We fix $\varepsilon > 0$ such that $u(z) < v(z) - 3\varepsilon$. Since u is continuous we can find a neighbourhood V of K such that $u > v - \varepsilon$ on V and since L is perfect we may assume that if $f_j(\zeta) \in \partial G$ then $f_j(\zeta) \in V$. For each j we consider domains $G_j = f_j^{-1}(G)$ which are unions of connected disjoint domains G_{jk} .

Taking averages if necessary we may assume that v is continuous so $u(z) < v(z) - \varepsilon$ on a ball $B = B(z, r)$. We denote by B_j sets $f_j^{-1}(B)$. The boundary Γ_{jk} of domains G_{jk} consists of points of U and points of S . We denote the first part by γ_{jk} , and the second by γ'_{jk} . If $u_j(\zeta) = u(f_j(\zeta))$ and $v_j(\zeta) = v(f_j(\zeta))$ then $u_j > v_j - \varepsilon$ on all γ_{jk} . Since $\text{supp } L \subset \partial D$ harmonic measures $a_j = \omega(0, \gamma'_{jk}, U)$ converge to 0. If E_j is the set of points in B_j where $\omega(\zeta, \gamma'_{jk}, G_{jk}) > s > 0$ then, evidently,

$$\omega(0, E_j, U) < \frac{a_j}{s}.$$

Let us introduce harmonic functions h_j on U with boundary values of u_j on S . Since $h_j \geq u_j$ on U the function

$$h_j \geq v_j - \frac{Aa_j}{s} - \varepsilon$$

on the set $C_j = B_j \setminus E_j$ and, therefore,

$$h_j > u_j - \frac{Aa_j}{s} + 2\varepsilon > u_j + \varepsilon$$

on C_j when j is sufficiently big. Since the leaf L is perfect, harmonic measures

$$\omega(0, B_j, U) > a > 0$$

and, therefore,

$$\omega(0, C_j, U) > a - \frac{a_j}{s} > \frac{a}{2}$$

when j is sufficiently big. So if we consider the universal holomorphic covering mappings g_j of $U \setminus C_j$ by U , and if we take mappings $q_j = f_j \circ g_j$ then

$$u(z_0) \leq \Phi(q_j) < \Phi(f_j) - \frac{a\varepsilon}{2},$$

and we get the contradiction.

Now if v is plurisuperharmonic and $v \geq u$ on K then, as before, we may assume that v is continuous on H . Repeating estimates from the first part of the theorem we may prove that for every $r > 0$ and for every $\varepsilon > 0$ the harmonic measure of points in B_j where $v_j \geq u_j$ is uniformly greater than 0 in U . For us it is enough to know that this set is non-empty because the combination of continuity of u and v and arbitrariness of r and ε implies that $v(z) \geq u(z)$. $\square$

The combination of Theorem 6.1 and 6.2 gives us

Theorem 6.3. *A continuous plurisubharmonic function u on a strongly pseudoconvex domain D in $\mathbb{C}^n$ is maximal if and only if through every point $z \in D$ passes the cluster of a perfect leaf L such that:*

- (1) u is maximal on $\text{cl } L$;
- (2) $\text{supp } L \subset \partial L$.

REFERENCES

1. E. Bedford, M. Kalka, *Foliations and complex Monge–Ampère equation*, Comm. Pure and Appl. Math. **30** (1977), 543–571.
2. H. Bremermann, *On a generalized Dirichlet problem for plurisubharmonic functions and pseudoconvex domains*, Trans. Amer. Math. Soc. **91** (1959), 246–276.
3. S. Bu, W. Schahermayer, *Approximation of Jensen measures by image measures under holomorphic functions and applications*, Trans. Amer. Math. Soc. **331** (1992), 585–608.
4. E. F. Collingwood, A.J. Lowater, *The Theory of Cluster Sets*, Cambridge University Press, 1966.
5. N. Danford, J. T. Schwartz, *Linear operators, part I: General Theory*, Interscience publishers, New York, London, 1958.
6. H. Goldmann, *An analytic algebra without analytic structure in the spectrum*, Ark. Math. **27** (1989), 89–95.
7. E.A. Poletsky, *To the Bloch theorem*, Uspekhi Mat. Nauk **41** (1986), 207–208.
8. E.A. Poletsky, *Plurisubharmonic functions as solutions of variational problems*, Proc. Symp. Pure Math., vol. 52, Part 1, 1991, pp. 163–171.
9. E.A. Poletsky, *Holomorphic currents (to appear)*.
10. W. Rudin, *Function theory in the Unit Ball in $\mathbb{C}^n$* , Springer-Verlag, Berlin Heidelberg New York, 1980.
11. A. Sadullaev, *Extensions of plurisubharmonic functions from submanifolds*, Dokl. Acad. Nauk UzSSR **5** (1982), 3–4.
12. G. Stolzenberg, *A hull with no analytic structure*, Jour. of Math. and Mech. **12** (1963).
13. J. Wermer, *Polynomial approximations on an arc in $\mathbb{C}^3$* , Ann. Math. **62** (1955), 269–270.
14. J. Wermer, *Polygonally convex hulls and analyticity*, Ark. Mat. **20** (1982), 129–135.

DEPARTMENT OF MATHEMATICS, 215 CARNEGIE HALL, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244